

FARAGO, Gyorgy, HA 5 BG

Untuned balancer and impedance transformer for short waves.
Radiotechnika 13 no.7:262-263 J1 '63.

FARAGO, Gyorgy (HA 5 BG)

Ultrashort wave transmission lines. Radiotechnika 13 no.9:332-333 S '63.

FARAGO, Gyorgy (HA 5 BG)

A significant meeting of radio amateurs in Gottwaldov. Radiotechnika
13 no.9:336-337 S '63.

PARAGO, George (HA 5 BG)

Supply lines in shortwave transmission technique. Radiotelegraph:
13 no. 10:372-373 0 '63.

FARAGO, Gyorgy (HA 5 BG) I. osztalyu rovidhullamu amator

Personal QSO with the Dresden amateur radio operators. Radio-
technika 13 no.11:414-415 N '63.

FARAGO, Gyorgy, (HA 5 BG)

Feeders in the shortwave transmission. Radiotechnika 13 no.12:
454-455 D '63.

FARAGO, Gyorgy, (HA 5 BG)

Problems relating to modern amateur transmitters. Pt.1.
Radiotechnika 14 no.2: 54-56 F'64.

FARAGO, Gyorgy (HA 5 BG) I. oszt.minositesu robidhullamu amator

Coil assembly made in the German Democratic Republic for
shortwave band receivers. Radiotechnika 14 no. 3:94-95
Mr '64.

Problems of modern amateur transmitters. Pt. 2. Ibid.:96-97.

PAVLOV, Gyorgy. HA 5 BG

Problems relating to modern amateur radio transmitters. Pt. 3.
Radiotekhnika 14 no. 4:134-135 Ap '64.

FARAGO, Gyorgy, (HA 5 BG)

Problems of modern radio transmitters. Pt. 4. Radiotechnika
14 no. 5:170-172 My '64.

FARAGO, Gyorgy (HA 5 BG), I. oszt. minositesu rovidhullamu amator

Problems of modern amateur transmitters. Pt. 5. Radiotechnika
14 no. 6:212-214 Je '64.

FABIAN, George, (UA 5 BG)

Problems of modern amateur transmitters. It.6. Radiotekhnika 14,
no.7:252-253 J1 '64.

HIDVEGI, Tibor; FARAGO, Gyorgy (HA 5 BG)

For modernizing amateur transmitters. Radiotechnika 15 no.7:
259-262 J1 '65.

1. Editorial Board Member, "Radiotechnika."

FARAGO, I.

On the origin and manifestations of narcotic diseases.
Magy. belorv. arch. 4 no.2:88-93 1951. (CML 20:11)

1. Doctor. 2. Neurological and Psychiatric Clinic (Director
Prof. Dr. Gyula Nyiro), Budapest Medical University.

FARAGO, I.

Syndrome of intervertebral disk hernia and multiple sclerosis. *Magy.*
belorv. arch. 5 no.3:133-136 Sept 1952. (CML 25:5)

1. Doctor. 2. Psychiatric and Neurological Clinic (Director -- Prof.
Dr. Gyula Myiro), Budapest Medical University.

FARAGO, I.

Marcomania, narcotic diseases. Orv. hetil. 93 no. 46:1312-1316
16 Nov 1952. (CIML 24:1)

1. Doctor. 2. Psychiatric and Neurological Clinic (Director -- Prof.
Dr. Gyula Nyiro), Budapest Medical University.

FARAGO, Istvan, dr.; KAZAR, Gyorgy, dr.

Connections of the early syndrome of skull injuries to later conditions; observations in 366 cases. Ideg. szemle 8 no.5: 152-156 Oct 55.

1. A Budapesti Orvostudományi Egyetem III. sz. Sebészeti Klinikája Traumatológiai Intézete (igazgató: Rubanyi Pál dr. egyetemi tanár) és Ideg Elme Klinikája (igazgató: Nyíró Gyula dr. egyetemi tanár) közleménye.

(HEAD, wounds & inj.

closed head inj., early & late neurotic & psychotic sympt. (Hun))

(WOUNDS AND INJURIES

head, closed inj., early & late neurotic & psychotic sympt. (Hun))

KULCSAR, Andor, dr.; MADOR, Gyorgy, dr.; ZOLCZER, Laszlo, dr.;
PARAGO, Istvan, dr.; MOLNAR, Edit, dr.

Clinical aspects and therapy of commotio cerebri. *Magy. Sebesszet* 10 no.1:4-13 Mar 57.

1. A Budapesti Orvostudományi Egyetem Baleseti Sebesszeti Intézetének Közleménye. Igazgató: Rubanyi, Pal, dr. egyetemi tanár.

(BRAIN, wounds & inj.

concussion, ther. & other clin. aspects (Hun))

FARAGC J										24									
H																			
<p>72. Condensers in refrigerating plants, by J. Faragó. ("Magyar Energetikaszemle" -- Power Economy in Hungary -- Vol. III, No. 7-12, pp. 143-146, July-Dec, 1950, 2 figs.)</p>										<p>two systems (assuming that the cooling tower has a cooling capacity of 100,000 cal) it is shown that the tower condenser requires 12 per cent more energy for the actual cooling besides necessitating the circulation of considerable additional amounts of water, which, in turn, increases the work of the water circulating pump. Applied to the above example, the cooling effect due to evaporation in the tower is at best less than 3,000 cal/hr, that is, 0.75 per cent of the 400,000 cal cooling capacity. On the other hand, it can be stated, however, that the use of the tower condenser is justified where cooling water is abundant, energy cheap, and where investment must be kept low.</p>									
<p>It was customary for domestic plants to erect the ammonia condensers of refrigerating plants in the open in the form of tower condensers, since their construction is less expensive than the horizontal baffled counterflow shell and tube condenser. Compared to the latter system, however, the temperature of condensation is higher and more pumping is required with tower condensers. On the basis of a practical example giving comparative figures for the</p>																			
<p>AS 8-51A METALLURGICAL LITERATURE CLASSIFICATION</p>																			

FARAGO, János, VI. oh.

Iodine reaction in pregnancy. Orv. hetil. 96 no.5:123-124 30
Jan 55.

1. A Fovarosí Tanács Egészségügyi Intézményei XI. Szakorvosi
Rendelőintézet (igazgató: Királyhegyi Robert dr.) Nőgyógyászati
Osztályának (főorvos: Pauncs Tivadar dr.) közleménye.

(IODINE,

test in pregn.)

(PREGNANCY TESTS,

iodine reaction)

FARAGO, Janos, dr.

Phenothiazines in combined obstetrical analgesia. Magy. noorv. lap.
25 no.2:102-105 Mr '62.

1. FM. Korvin Otto korhaz szuleszeti-nogyogyaszati osztalyanak
(Foorvos: Fabian Laszlo dr.) kozlemenye.

(PHENOTHIAZINES ther) (ANESTHESIA OBSTETRICAL)

1ST AND 2ND ORDERS										3RD AND 4TH ORDERS									
<p>FARAGO, K.</p> <p>777</p> <p>7</p> <p>*Normal Potentials of Lead and Silver. Károly Faragó (<i>Acta Lit. Sci. Regiae Univ. Hung. Franciscus-Josephinus, Sect. Chem., Min., Phys.</i>, 1937, 8, 193 210; <i>C. Ab.</i>, 1938, 32, 2153).—A modified form of the Huggenheim glass electrode made of Jena glass was used. The normal potential of the lead electrode, determined in solutions of $Pb(NO_3)_2$, $E_0 = 0.1274$ v., agrees with the value calculated by Frumkin. The activities and activity coeffs. of the various lead ions are calculated and tabulated. A suitable silver electrode was made of a metallic silver plate in a glass tube, one end of which was closed with paraffin and then filled with paraffin. Electrolytic silver was then deposited on this electrode covered with parchment paper to prevent precipitation of possible impurities. This electrode should be placed in the solution to be studied some hours before measurement, and hydrogen should be led through the solution during the determination. The normal potential found for silver in perchlorate solutions, $E_0 = 0.7905$ v., agrees with the value of Lewis and Randall. —N. (7).</p>																			
<p>ASB-SLA METALLURGICAL LITERATURE CLASSIFICATION</p> <p>FROM 834179</p>																			

FARAGO, Kalman

Data on the question of planning residential area units. Magyar
ipar 12 no.8:379-383 '63.

FARAGO, Katlin, dr.; HAJOS, Endre, dr.

Simultaneous tomography. Magy. radiol. 8 no.2:104-106 May 56.

1. A Budapesti Orvostudományi Egyetem Röntgenklinikájának
(Igazgató: Ratkoczy, Mándor, dr. egyet. tanár) közleménye.
(ROENTGENOGRAPHY
tomography, simultaneous of several layers, new
technic (Hun))

FARAGO, Katalin, dr.; GIMES, Bela, dr.

Radiotherapy of myasthenia gravis pseudoparalytica. Orv.hetil.
100 no.48:1732-1734 N '59.

1. A Budapesti Orvostudományi Egyetem Röntgenklinikájának
(igazgató: Ratkoczy Mándor dr. egyetemi tanár) közleménye.
(MYASTHENIA GRAVIS radiother)

FARAGO, L.

Significance of sternal puncture in dermatology. *Borogyog vener. szemle*
5 no.6:168-171 Dec 51. (CML 21:4)

1. Doctor. 2. Skin and Venereal Diseases Clinic (Director—Prof. Dr.
Ferenc Foldvari), Budapest University.

KAJTOR, Ferenc; FARAGO, Lajos; TOROK, Pal

Effect of peripheral sensory stimulation on the convulsive activity of the hippocampus in evipan anesthesia. Ideg szemle 10 no.5-6:171-180 Oct-Dec 57.

1. A Debreceni Orvostudományi Egyetem Ideg-elmeklinikájának közleménye.
(HIPPOCAMPUS, in various dis.
epilepsy, temporal, EEG of hippocampal responses to sensory stimulation of peripheral nerves in hexobarbital anesth. (Hun))
(EPILEPSY, physiol.
EEG of hippocampal responses to sensory stimulation of peripheral nerves in hexobarbital anesth. in temporal epilepsy (Hun))
(ELECTROENCEPHALOGRAPHY, in various dis.
epilepsy, temporal, EEG of hippocampal responses to sensory stimulation of peripheral nerves in hexobarbital anesth. (Hun))
(NERVES, PERIPHERAL, in various dis.
same)

HULLAY, Jozsef; FARAGO, Lajos; TOROK, Pal

Data on ~~memory~~ functions based on electric cortical stimulations in temporal epilepsy. Ideg. szemle 11 no.1-2:15-17 Feb-Apr 58.

(MEMORY,

funct. of memory cortex studies by electric cortical stimulations in temporal epilepsy (Hun))

(~~CEREBRAL CORTIX~~, physiol.

memory cortex funct. studied by electric cortical stimulations in temporal epilepsy (Hun))

EXCERPTA MEDICA Sec 13/Vol 13/1 Dermatology Jan 59

210. KAPOSI'S SARCOMA - Sarcome de Kaposi - Farago L. Budapest -
ANN. DERM. SYPH. (Paris) 1958, 85/3 (297-298) Illus. 2

Case report.

(XII, 5, 16)

FARAGO, Laszlo, dr.

"The process of thinking and the regularities of analysis, synthesis, and generalization," edited by S.L. Rubinshteyn. Reviewed by Laszlo Farago. Magyar pszichológiai szemle 18 no.3:374-376 '61.

FARAGO, Laszlo, tanszekvezeto

International symposium on teaching mathematics held in Budapest.
Magy tud 69 no.12:784-788 D '62.

1. Orszagos Pedagogiai Intezet.

FARAGO, Laszlo, dr.

Psychological problems relating to the up-to-date teaching of mathematics. Magyar pszichol szemle 20 no.3:441-448 '63.

1. Orszagos Pedagogiai Intezet.

HUNGARY

FARAGO, Dr. Laszlo, and HOLCZINGER, Dr. Laszlo, Ear-Nose-Throat Department (Orr-Gege-Fulosztaly) of the National Institute of Oncology (Orszagos Onkologiai Intezet) and Research Institute for Oncopathology (Onkopathological Research Institute).

"Tumors of the Salivary Glands"

Budapest, Magyar Onkologia, Vol 10, No 4, Dec 1966; pp 236-239.

Abstract [Authors' English summary, modified]: After subjecting to pathological and clinical consideration tumors originating from the salivary glands of the oral cavity, authors discuss 17 of their own cases diagnosed and treated during 1956-1965. Their cases were mostly localized in the palate. Eleven cases of pleomorphic adenomas, 1 pleomorphic adenoma with malignant development, 4 adenocarcinomas, 1 mucoepidermoid tumor were treated. Combined therapy is considered to be the most effective method of treatment. Of the 12 patients with pleomorphic adenoma, 9 survived for more than 5 years, 2 survived for more than 3 years, 1 died after having been free of symptoms for two years; of the 4 cases of adenocarcinoma there were 3 relapses, including one after 9 years; 2 are alive and 2 died. A patient with mucoepidermoid tumor has been free of symptoms for two years. 41 References, mainly Western.

FARAGO, M.

FARAGO, M. - The use of a color filter in microphotography. p. 90.
Vol. 2, no. 4, Aug. 1956
Képes Hangtechnika. Budapest, Hungary

SOURCE: East European Accessions List (EEAL) Vol. 6, No. 4--April 1957

FARAGO, M.

The cell wall. p. 8. (Magyar Mezogazdasag, Vol. 11, no. 3, Feb. 1956 Budapest)

SO: Monthly List of East European Accession (EEAL) LC, Vol. 6, no. 7, July 1957. Uncl.

FARAGO, M.

FARAGO, M. Plante resistant to freezing. p. 11

Vol. 11, no. 8, A pr. 1956

MAGYAR MEZOGAZDASAG

AGRICULTURE

Budapest, Hungary

So: East European Accession, Vol. 6, No. 3, March 1957

FARAGO, Mihaly, dr. (Fertod); HORVATH, Karoly (Pecs, Alkotmany u.20);
KINDLER, Andras

Miscellany. Radiotechnika 11 no.11:341 N '61.

1. TV muszeresz, Orion Radio es Televizio Szerviz (for Kindler).

116

FARAGO, Pal
ca

PROCESSES AND PROPERTIES INDEX

The histamine treatment of allergic patients. *Pal Farago, Orvosi Hetilap 81, 945-6 (1937).*—Clinical exper. Since 1933 showed that doses of histamine beginning at 0.001-0.01 mg. and increasing every 2nd day usually have good effects. The healing effect probably is due to the acclimatization of the organism to free histamine.
S. S. de Pinálv

ASB-51A METALLURGICAL LITERATURE CLASSIFICATION

180000 181000 182000 183000 184000 185000 186000 187000 188000 189000

180000 181000 182000 183000 184000 185000 186000 187000 188000 189000

1		2		3		4		5		6		7		8		9		10		11		12		13		14		15		16		17		18		19		20		21		22		23		24		25		26		27		28		29		30		31		32		33		34		35		36		37		38		39		40		41		42		43		44		45		46		47		48		49		50		51		52		53		54		55		56		57		58		59		60		61		62		63		64		65		66		67		68		69		70		71		72		73		74		75		76		77		78		79		80		81		82		83		84		85		86		87		88		89		90		91		92		93		94		95		96		97		98		99		100		101		102		103		104		105		106		107		108		109		110		111		112		113		114		115		116		117		118		119		120		121		122		123		124		125		126		127		128		129		130		131		132		133		134		135		136		137		138		139		140		141		142		143		144		145		146		147		148		149		150		151		152		153		154		155		156		157		158		159		160		161		162		163		164		165		166		167		168		169		170		171		172		173		174		175		176		177		178		179		180		181		182		183		184		185		186		187		188		189		190		191		192		193		194		195		196		197		198		199		200		201		202		203		204		205		206		207		208		209		210		211		212		213		214		215		216		217		218		219		220		221		222		223		224		225		226		227		228		229		230		231		232		233		234		235		236		237		238		239		240		241		242		243		244		245		246		247		248		249		250		251		252		253		254		255		256		257		258		259		260		261		262		263		264		265		266		267		268		269		270		271		272		273		274		275		276		277		278		279		280		281		282		283		284		285		286		287		288		289		290		291		292		293		294		295		296		297		298		299		300		301		302		303		304		305		306		307		308		309		310		311		312		313		314		315		316		317		318		319		320		321		322		323		324		325		326		327		328		329		330		331		332		333		334		335		336		337		338		339		340		341		342		343		344		345		346		347		348		349		350		351		352		353		354		355		356		357		358		359		360		361		362		363		364		365		366		367		368		369		370		371		372		373		374		375		376		377		378		379</	
---	--	---	--	---	--	---	--	---	--	---	--	---	--	---	--	---	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-----	--	-------	--

APC
5-118

Doc. 11111

328. Electron multiplier tube of large effective cathode surface area. by
P. S. Farago. *Nature*, 161. p. 60. January 10. 1948.

Methods for increasing the effective surface of electron multiplier tubes for
particle counting are described. The effective surface may be increased by
projecting the electron-optical image of a cathode of large surface area on the
first multiplying electrode of a multiplier tube of the usual dimensions or by
increasing the length of all electrodes of a circular-type multiplier. The
development of the tube for the purpose of researches on cosmic radiation is go-
ing on.

SA		<p>1408. Measurement of extremely small light intensities. P. S. FARAGO AND E. F. POCEA. <i>Publ. Univ. Tech. Sci., Budapest</i> (No. 1) 10-14 (1949).</p> <p>Using a phototube with a dark current of 10^{-11} A. photocurrents $\sim 10^{-14}$ A can be detected by modulating the incident light with a suitable frequency (50, 120 and 240 c/s were used) and amplifying the resultant total current by a tuned circuit with bandwidth 1-3 c/s. For d.c. operation, the dark current of the cell may be balanced out with sufficient accuracy if very stable sources of supply are used.</p> <p>M. CORCORAN</p>		<p>535.247.4</p> <p>AS3</p> <p>J</p>	
A10.31A METALLURGICAL LITERATURE CLASSIFICATION		1000 17101194		1000 17101194	
100000		100000		100000	

CA

3

Methods for measuring small light intensities. P. S. Farago. *Hung. J. Phys.* 1, No. 6, 913 (1949) (in English). - In the methods used, either the usual photocell is replaced by a multiplier phototube and the continuous photocurrent is measured after amplifying by electron tubes, or the current is measured after amplifying by counting the no. of discrete single photons emitted by the cathode of the multiplier tube. For a given multiplier tube the limit of both methods is quantitatively the same, if the frequency band width in the former case, or if the period of counting in the latter case is properly chosen. The selective amplifier method has the advantage that by using 2 channels of the

same band width, one tuned to the light modulation frequency, the other to a different one, photocurrent and noise can be detd. simultaneously. Multiplier phototubes with very small dark emission at room temp. were produced with the aid of photocathodes and multiplying dynodes which have a higher work function than that of the Ag-Cs-Cs layer. Although the efficiency of the cathode is thereby somewhat reduced, such a multiplier tube was more suitable for measuring extremely small light intensities, especially in the blue region. Jovanovic

1951

3

CA

Probability distribution of the number of secondary electrons. P. S. Farago and L. Takacs (Univ. Tech. Sci.,

Budapest, Hung.). *Hung. Acta Phys.* 1, No. 6, 43-52 (1949) (in English). -The probability distribution of the pulse amplitudes produced by single electrons in a $(k-1)$ - and a k -stage electron multiplier tube is discussed, and a simple relation is derived for the generating functions of these distributions. Based on this relation a suitable method is given for the detn. of the probability distribution of the no. of secondary electrons released by 1 electron from a secondary emitting surface. The soln. of the problem is extended to the case of arbitrary primary particles. A special method is given for the detn. of the efficiency of multiplier tubes as electron counters. István Földi

1957

SIGNATURE																										DATE																									
SIGNATURE													DATE													SIGNATURE													DATE												
SA																										A53																									
335,247.4																																																			
7100. Notes on some methods for measuring small light intensities. P. B. FARACH. <i>Revue de Phys.</i> , 1 (No. 6) 9-13 (1958).																																																			
Discuss the 2 methods commonly used for measuring very small light intensities with a multiplier photocell, viz. (i) direct measurement of the photocurrent due to continuous exposure, and (ii) modulation of the incident beam at some convenient frequency. The most favorable method depends on the conditions met with, particularly in regard to light intensity to be measured. J. V. T. WALSH																																																			
AND SEA BIBLIOGRAPHICAL LITERATURE CLASSIFICATION																																																			
SIGNATURE																										DATE																									
SIGNATURE																										DATE																									

Hungary/Radiophysics - Superhigh Frequencies, I-11

Abst Journal: Referat Zhur - Fizika, No 12, 1956, 35391

Author: Farago, Peter; Groma, Geza

Institution: None

Title: Reflex Klystrons

Original

Periodical: Magyar tud. akad. Kozl. fiz. kutato intez. kozl. 1953, 1, No 1-2,
66-81; Hungarian

Abstract: See Referat Zhur - Fizika, 1955, 9786

Card 1/1

FARAGO, P. S.

HUNG.

10423. On the application of a simple magnetic resonance absorption device for quantitative measurements. P. S. FARAGO, M. GILES AND J. MERTZ - *Ann. Phys. Hongk.*, 3, No. 3-4, 139-151 (1954).

A specially simple nuclear resonance detection is obtained if the sample is in the resonant circuit of an oscillator and the power output is observed as a function of magnetic field. The linearity of change of output with γ is demonstrated.

FARAGO, P.S.

5

621.385.1.029.6 : 621.373.423
5221. Reflex oscillators. P. S. FARAGO AND G.
OROM. *Acta phys. Hungar.* **4**, No. 1, 7-22 (1954).
The mechanisms of reflex klystron and Barkhausen-Kurz oscillators were compared and found to be essentially similar. Both are feedback oscillators, with one h.f. field producing the bunching of electrons and also converting (a fraction of) the kinetic energy of the electrons into h.f. power, with a feedback mechanism maintained by a constant retarding field. All the differences in the behaviour of the two types of oscillators can be explained by the fact that in the reflex klystron, the h.f. interaction space is separated from the constant retarding field, and in the Barkhausen-Kurz oscillator the two are superimposed. The main consequences are discussed quantitatively.

PT
JLH

FARAGO, P.S.

4

HUNG

530.145 : 537.2

10787. Quantum effects in the interaction between
free electrons and electromagnetic fields. P. S.
FARAGO AND G. MARX. *Acta phys. Hungar.*, 4,
No. 1, 23-30 (1954).

An electron beam shot through a transverse r.f.
field suffers a direction spread. If experimental
conditions are suitably chosen, the direction spread
seems to be due only to the quantum dispersion of
energy exchange between free electrons and r.f. field.
A simple collector electrode system might allow not
only the detection of the direction spread of the
electrons, but the presence of a quantum effect might
be checked by plotting the collector current versus r.f.
field amplitude, the plot for the quantum effect being
different from those for classical effects.

RDW

FARAGO, P.S.

2879

INVESTIGATION OF MAGNETIC MOMENTS OF ATOMIC
NUCLEI. P. S. Farago, M. Gács, and J. Mertz (Central
Research Inst. of Physics, Budapest). Nuovo cimento
(10) 2, 1110-13 (1955) Nov. *NU*

Measurements of the nuclear magnetic moments of Na^{23} ,
 P^{31} , and F^{19} were repeated. The results are summarized
and compared with those of other authors. (B.J.H.) *(2)*

BMF

PM

FARAGO, P. S.

V 7216

PH

QUANTUM EFFECTS IN THE INTERACTION BETWEEN
FREE ELECTRONS AND ELECTROMAGNETIC FIELDS.
P. S. Farago (Central Research Inst. for Physics, Budapest,
Hungary) and G. Marx (Physical Inst. of the Eotvos Univ.,
Budapest, Hungary). Phys. Rev. 99, 1063-4(1955) Aug. 15.

An electron beam shot through a transverse rf field may suffer a directional spread. If experimental conditions are suitably chosen, the directional spread may be due only to the quantum dispersion of energy exchange between free electrons and rf field. A simple collector electrode system might allow not only the detection of the directional spread of the electrons, but the presence of a quantum effect might be checked by plotting the collector current versus rf field amplitude, the plot for the quantum effect being different from those for classical effects. The results of various theoretical treatments of the effect are briefly compared, both from the point of view of their principal foundations and of the possibility of their experimental verification. (auth)

Small 1
RXX

FARAGO, Peter

HUNGARY/Electronics - General.

Abs J. r : Ref Zhur - Fizika, No 6, 1959, 13444

Author : Farago, Peter

Inst : -

Title : Concerning on the Book by V.F. Vlasov "Electric Vacuum Devices"

Orig Pub : Magyar tud. akad. Mat. es fiz. tud. OSZT. kozl., 1956, 6, No 2, 262-264

Abstract : The book is a translation of the second Russian edition (1949) and is used as a text for higher institutions of learning. The first 22 chapters have material of the course of lectures on electric vacuum devices, delivered by the author in the electrotechnical higher technical institutions for communication.

Card 1/1

P. S. FARAGO

4

Review of the experimental evidence for the law of variation of the electron mass with velocity. P. S. Farago and L. Jancsó (Central Research Inst. Phys., Budapest, Hung.). *Nuovo Cimento* (10), 8, 1411-30 (1967) (in English).—P. and

TH
UP
review the exptl. measurements which support the Lorentz equation for the variation of mass with velocity. Although all expts. are consistent with this law, they conclude that only the data on the fine structure of H-like spectra (Ottacher, C.A. 12, 1147) are sufficiently precise to distinguish between the Lorentz equation and that of Abraham, derived for a rigid electron (no reference given).

George L. Cunningham

gr.

AUTHORS: Zrelov, V. P., Tyapkin, A. A., Farago, P. S. SOV/56-34-3-4/55

TITLE: Measurement of the Proton Mass at 660 MeV (Izmereniye massy protonov pri energii 660 MeV)

PERIODICAL: Zhurnal Eksperimental'noy i Teoreticheskoy Fiziki, 1958, Vol. 34, Nr 3, pp. 555-558 (USSR)

ABSTRACT: The present work compares the values computed by means of the relativistic relation $m_2 = m_0 [1 - (v^2/c^2)]^{-1/2}$ based on the velocity measured with the values $m_1 = p/v$ of the mass which were determined from the measured momenta and velocities of protons. The measurements were made on an external proton beam with about 660 MeV which made essentially easier the determination of possible errors. The general scheme of the measuring device is shown in a diagram. The external beam of a 6 m-synchrocyclotron passes a system of collimators, then was deflected within the field of an electromagnet with a pole diameter of 1 m, passed a second collimator and then impinged upon ionization chamber. The control measurements are also described. In the determination of the momentum of protons by means of a current carrying conductor the values

Card 1/3

Measurement of the Mass of 660 MeV Protons

SOV/56-34-3-4/55 . . .

$Q = 300.0 \pm 0.3$ and $I = 0.681 \pm 0.001$ amp. were obtained. Q denoting the load applied to a silk thread. From this the value $p = 1296.5 \pm 2.3$ MeV/c is obtained for the momentum. Various measurements carried out at $Q = 200,0$ g showed results which coincide within the limit of measuring errors with earlier obtained results. Various details of the measurements are discussed. Also the second correction of the energy loss in air must be taken into account which amounts to $\Delta E_2 = 1.5$ MeV. The total correction of the energy amounts to $\Delta E = \Delta E_1 + \Delta E_2 = 8.1$ MeV. The authors intend to determine the deviations from the fundamental law of relativistic theory $m = m_0 [1 - (v^2/c^2)]^{1/2}$, and use relativistic relations in the determination of the corrections ΔE and Δv . When the found values for the momentum and velocity of the protons are taken into account $m_1 = p/v = 1598.2 \pm 3$ MeV/c² and $m_2 = m_0 [1 - (v^2/c^2)]^{-1/2} = 1604.3 \pm 1.9$ MeV/c² are obtained. From this further results $\Delta m = m_2 - m_1 = 6.1 (1 \pm 0.5)$ or $\Delta m/m = 0.004 (1 \pm 0.5)$. The errors mentioned are the mean square deviations. Thus the results obtained here coincide with the relativistic law for the increase of mass with increasing velocity within the error limits mentioned.

Card 2/3

Measurement of the Proton Mass at 660 MeV

SOV/56-34-3-4/55

There are 1 figure and 9 references, 2 of which are Soviet.

ASSOCIATION: Ob"yedinennyy institut yadernykh issledovaniy
(United Institute for Nuclear Research)

SUBMITTED: September 12, 1957

Card 3/3

PATAKI, Pal, dr.; PANGSOVAY, Jozsef, dr.; FARAGO, Peter, dr.

Neurinoma of the extremity. Orv.hetil. 102 no.4:174-175 22 Ja'61.

1. Fovarosí Arpad Kórház, Sebészeti Osztály.
(NEUROMA case reports)
(EXTREMITIES neopl)

FARAGO, Stefan

. Planning of train service and its performance. Zel dop tech
12 no.9:241-242 '64.

1. Dispatcher, SVD.

47531-66 EWR J: IM
ACC NR: AT6035004

SOURCE CODE: HU/2502/66/047/002/0173/0184

AUTHOR: Szoke, Sandor--Seke, Sh. (Doctor) and Farago, Terez, of the Central Research Institute for Chemistry at the Hungarian Academy of Sciences in Budapest.

"Calculation of the Force Field on the Ethylene Model"

Budapest, Acta Chimica Academiae Scientiarum Hungaricae, Vol 47, No 2, 1966, pp 173-184.

Abstract: [English article; authors' English summary] The force constants of the ethylene molecule have been calculated. Fourteen equations were used to calculate eleven force constant components applying the method of least squares. In the calculation of the constants of the stretching vibrations, the force constants of diatomic molecules were used as approximate values. By this procedure the relationship of the force constants with the applied empirical data has been established.

16 formulas and 6 tables. APRS: 36,002

Orig. art. has: 2 figures,

TOPIC TAGS: ethylene, molecular physics

SUB CODE: 07,20 / SUBM DATE: 22 Jul 65 / ORIG REF: 001 / OTH REF: 019
SOV REF: 004

Card 1/1

0921 1577

FAK 050, T.

200

Faragó, T. Über das arithmetisch-geometrische Mittel.

Publ. Math. Debrecen 2, 150-156 (1951).

The arithmetic-geometric mean of two positive numbers a_0, b_0 is the common limit $M(a_0, b_0)$ of the two sequences $\{a_n\}, \{b_n\}$ generated by $a_{n+1} = \frac{1}{2}(a_n + b_n)$, $b_{n+1} = (a_n b_n)^{1/2} > 0$. For these sequences, we have

$$(*) \quad b_n \leq b_{n+1} \leq M(a_0, b_0) \leq a_{n+1} \leq a_n.$$

For complex numbers a_0, b_0 , the author now considers the same sequences, except that now the condition $b_{n+1} > 0$ is generalized to the restriction that b_{n+1} lie in the angle $\alpha_{n+1} = \frac{1}{2}(\alpha_n, 0, \beta_n)$, with $0 \leq \alpha_n \leq \pi$. Clearly we do not now necessarily have $|b_n| \leq |a_n|$; nevertheless, the following analogue of (*) is given:

$$|b_n| \cos \frac{1}{2}\alpha_n \leq |b_{n+1}| \cos \frac{1}{2}\alpha_{n+1} \\ \leq |M(a_0, b_0)| \leq \frac{1}{2}(|a_n| + |b_n|) \leq \frac{1}{2}(|a_{n+1}| + |b_{n+1}|).$$

Also, necessary and sufficient conditions on (a_0, b_0) and (a'_0, b'_0) , in order that $|M(a_0, b_0)| \leq |M(a'_0, b'_0)|$, are given. E. F. Beckenbach (Los Angeles, Calif.).

Source: Mathematical Reviews,

Vol. 13 No. 10

~~SECRET~~ FARAGO, TIBOR

Farago, Tibor. Contribution to the definition of group.
~~Publ. Math. Debrecen 3 (1953), 133-137 (1954).~~

Let G be a system of elements in which a single-valued binary multiplication is defined. Suppose G contains a unit element and that every element of G has a left inverse. If, in addition, G is associative, then G is a group. The author considers fifteen substitutes for the associative law, which are obtained by changing the position of the brackets and/or permuting the three elements a , b , and c in the right member of $(ab)c = a(bc)$. The associative law in G is replaced in turn by each of these fifteen substitutes. Nine of the resulting systems are commutative groups, three others are commutative systems not necessarily groups, while the remaining three (one of which is necessarily a quasigroup) need be neither commutative or associative. *D. C. Murdoch.*

Distb: hF1

1. Determination of higher-order partial derivatives of the velocity function from known values along a stream surface. (In English) T. PARÁK. Acta Technica Academiae Scientiarum Hungaricae. Vol. 20, 1958, No. 3-4, pp. 217-227.

The formula of the n th derivatives is deduced and verified assuming the velocity distribution on the stream surface $r(u, v)$ to be known. The n th partial derivatives in a system of cylindrical co-ordinates are also given. Certain principles concerning the analytical specification of velocity components on known stream surfaces are established.

FARAGO, T. (Budapest, XI., Sztoczek u.2-4)

An elementary method for an approximate determination of the
real roots of algebraic equations. Periodica polytechn electr
7 no. 3:197-208 '63.

1. Lehrstuhl für Mathematik Techn. Univ. Budapest.
" abgelegt von Dr.E.Makai.

FARAGO, Tibor, dr. (Budapest, XI., Sztoczek u.2-4)

Studies on polynomials with one or more variables. Periodica
polytechn electr 8 no.2:193-206 '64.

1. Lehrstuhl für Mathematik, Technische Universität, Budapest.
Vorgelegt von Prof. Dr.I. Fenyő.

FAIDANO, V.I.

Rapid method of repairing open-hearth furnaces. Metallurg 19 no.7:
29 J1 '65. (MIRA 18:7)

1. Nachal'nik otдела tekhnicheskoy informatsii Taganrogskogo
metallurgicheskogo zavoda.

FARAKHOV, A.A.

"Correction of Off-shore oil Extraction Structures and Catode protection."
(Korroziya Morskikh Neftepromyslovykh Sooruzhenii i Katodnya zashchita.)
Baku, Aznefteizdat, 1955.

PARAMAZOV, SAYRAN ARUTYULOVICH

GUSEYNOV, Dzhebrail Alekper ogly; PARAMAZOV, Sayran Arutyulovich; SPEKTOR,
Sh.Sh., red.; AL'TMAN, T.B., red. izd-va.

[Technology and mechanisation of the production of petroleum bitumen]
Tekhnologiya i mekhanizatsiya proizvodstva neftebitumov. Baku,
Azerbaidzhanskoe gos.izd-vo nef't.i nauchno-tekhn.lit-ry, 1957. 180 p.
(MIRA 11:1)

(Petroleum)

(Bitumen)

FARAMAZOV, S. A., Cand Tech Sci (diss) -- "The problem of mechanizing laborious processes in the production of high-melting bitumen". Baku, 1960. 10 pp
(Min Higher and Inter Spec Educ USSR, Azerb Order of Labor Red Banner Inst of Petroleum and Chem im M. Azizbekov), 150 copies (KL, No 12, 1960, 128)

FARAMAZOV, Seyran Arutyunovich; CHIZHOV, A.A., ved. red.;
DEM'YANENKO, V.I., tekhn. red.

[Complete mechanization and automation of the production
of solid oil asphalt] Kompleksnaia mekhanizatsiia i avto-
matizatsiia proizvodstva tverdykh neftiannykh bitumov. Le-
ningrad, Gostoptekhnizdat, 1963. 122 p. (MIRA 16:10)
(Asphalt)

FARAMAZOV, S.A.; ABDULOVA, Z.M.

Efficient positioning of return band chambers in pipestills.
Mash. i nef. obor. no.8:23-24 '64. (MIRA 17:11)

1. Bakinskiy nefepereperabatyvayushchiy zavod im. XXII s"yezda
Kommunisticheskoy partii Sovetskogo Soyuza.

FARAMAZOV, S.A.; KOSENKOV, V.G.; AKHMEDOV, K.R.

Stabilizing the draft of pipestill flues. Nefteper. i neftekhim.
no.4:45-47 '65. (MIRA 18:5)

1. Bakinskiy neftepererabatyvayushchiy zavod im. XXII s"yezda
Kommunisticheskoy partii Sovetskogo Soyuza.

FARMAZOV, S.A., kand. tekhn. nauk; GINZBURG, M.B., inzh.; PIRUMYAN, M.Ye.,
inzh.; TSOYREF, M.I., inzh.

Mechanization of the cutting of a high-viscosity polymer. Mekh.
i avtom. proizv. 19 no.10:11-12 0 '65. (MIRA 18:12)

KARAMYAN, K.A.; PARAMAZYAN, A.S.

Mineralization phases of the Kadsharan copper and molybdenum
deposit. Izv. AN Arm. SSR. Geol. i geog. nauki 13 no.3/4:65-88
'60. (MIRA 13:9)

1. Institut geologicheskikh nauk AN ArmSSR.
(Kadsharan region (Armenia)—Mineralogical chemistry)

MELIKSETYAN, B.M., PARAMAZYAN, A.S., KHURSHUDYAN, E.Kh.

Tellurobismuthite and certain other tellurides from the Kaler molybdenite deposit. Dokl.AN Arm.SSR 30 no.4:239-244 '60.

(MIRA 13:8)

1. Institut geologicheskikh nauk Akademii nauk ArmSSR. Predstavleno akad. AN Armyanskoy SSR I.G.Magak'yanom.
(Megri District--Tellurides)

S/081/61/000/022/015/076
B102/B108

AUTHOR: Faramazyan, A. S.

TITLE: Lawfulness of rhenium distribution in ores from
Kadzharan

PERIODICAL: Referativnyy zhurnal. Khimiya, no. 22, 1961, 97, abstract
22G108 (Izv. AN Arm.SSR. Geol. i geogr. n., v. 14, no. 1,
1961, 39-56)

TEXT: Re in molybdenites was determined colorimetrically according to the color of its rhodanide complex. Mo was previously separated by Hg(1+) nitrate or ascorbic acid. The Re contents in molybdenites in the following stages of mineralization are presented (in 10^{-2} %): 1) Quartz-magnetite (2.6-3.0), 2) quartz-field spar (3.4-4.1), 3) quartz-molybdenite (3.4-4.9, average 3.21), 4) quartz-chalcopyrite-molybdenum (3.0-16.1, average 7.39), 5) quartz-chalcopyrite (2.2-3.2, average 2.78), 6) quartz-pyrite (3.0-3.7, average 3.27). Just like molybdenite which is segregated mainly in the 3rd (70 %) and 4th (15 %) stages, also the main mass of Re is segregated in these stages: 61.2 and 30.2 %, respectively.

Card 1/2

Lawfulness of rhenium distribution...

S/081/61/000/022/015/076
B102/B108

The Se and Te contents were also determined in these molybdenites by semiquantitative spectroscopical and chemical methods. A diagram is given which shows that the Se content increases with increasing Re content. This is explained by a geochemically similar behavior of these elements. The Mo and Re contents of some other sulfides are also given; the Mo:Re-ratio is the same as in the molybdenites. This leads to the assumption that Re in sulfides is due to mechanical molybdenite admixtures. A comparison of the Mo:Re ratios in molybdenite (2000:1), in powellite (10⁵:1), and in mine water (200:1) shows that Re is removed from the oxidation zone. [Abstracter's note: Complete translation.]

✓

Card 2/2

FARAMAZYAN, A.S.; KHACHADURIAN, E.Kh.

Enargite and luzonite from the Kadzharan copper-molybdenite deposit.
Zap.Arm.otd.Vses.min.ob-va no.2:5-17 '63. (MIRA 16:10)

FARAMAZYAN, A.S.; AKOPYAN, A.G.

Rhenium and some molybdenum ore manifestations in the Ayotsdzor ore region. Izv. AN Arm. SSR. Geol. i geog. nauki 16 no. 3: 61-66 '63.

(MIRA 17:2)

1. Institut geologicheskikh nauk AN Armyanskoy SSR.

MAGAK'YAN, I.G., akademik; PIDZHYAN, G.O.; FARAMAZYAN, A.S.

Rhenium in copper-molybdenum deposits of the Armenian
S.S.R. Dokl. AN Arm. SSR 37 no.2:77-81 '63. (MIRA 17:2)

1. Institut geologicheskikh nauk AN Armyanskoy SSR.
Akademiya nauk Armyanskoy SSR (for Magak'yan).

FARAMAZYAN, A.S.; KHURSHUDYAN, E.Kh.

Isomorphism between molybdenum and rhenium in molybdenites.
Dokl. AN Arm. SSR 37 no.4:211-216 '63. (MIRA 17:8)

1. Institut geologicheskikh nauk AN ArmSSR. Predstavleno akademikom AN Armyanskoy SSR I.G. Magak'yanom.

FARMAZYAN, R.

American "aid" and Turkish finances. Fin.SSSR 20 no.8:
81-89 Ag '59. (MIRA 12:11)
(Turkey--Economic assistance--American)
(Turkey--Finance)

FARAMAZYAN, Rachik Artashesovich; SAMYKIN, S., red.; RODIONOVA, L.,
Mlad. red.; NOGINA, N., tekhn. red.

[Economy of present-day Canada] Ekonomika sovremennoi Kanady.
Moskva, Sotsekgiz, 1963. 222 p. (MIRA 16:10)
(Canada--Economic conditions)

KILESSO, A.I.; FARAMAZYAN, R.A.; KONONYUK, B.Z.; MARTINSEN, Z.A.;
ANDREYEV, Yu.V.; SLAVIN, S.V.; RUSETSKIY, S.B.; GLUSHKOV,
V.P., otv. red.; PLISKINA, Ye.M., red.; TIKHOMIROVA, S.G.,
tekhn. red.

[The shipbuilding industry of capitalist countries] Sudostroitel'naya promyshlennost' kapitalisticheskikh stran.
Moskva, Izd-vo AN SSSR, 1963. 471 p. (MIRA 16:10)

1. Akademiya nauk SSSR. Institut mirovoy ekonomiki i mezhdunarodnykh otnosheniy.
(Shipbuilding)

FARAN, Antonin, inz.

Automatic recorder in railroad stations. Zel dop tech 11
no.3:85-86 '63.

PARANA, J.

Wstępna kontrola gospodarki materiałowej (Preliminary control of material economy), by J. Parana. Reported in New Epoch, (Nowe Zisaki), No. 6, March 15, 1956.

FARAPONOVA, E.A.

Meeting on psychological problems of industrial education and training in the school. Vop.psikhol. 7 no.2:188-189 Mr-Ap '61.
(MIRA 14:6)

(Work)

ZAPOROZHETS, A.V. (Moskva); FARAFONOVA, E.A. (Moskva)

Work of the Moscow Section of the Society of Psychologists in
1960-1964. Vop. psikhol. 11 no.2:189-190 Mr-Ap '65.
(MIRA 18:6)

FARAPONOVA, G.P.

Some results of actinometric observations from free aerostats.
Trudy TSAO no.8:27-35 '52. (MIRA 12:1)
(Aeronautics in meteorology) (Solar radiation)

124-58-9-10068

Translation from: Referativnyy zhurnal, Mekhanika, 1958, Nr 9. p 90 (USSR)

AUTHORS: Faraponova, G. P., Kastrov, V. G.

TITLE: Actinometric Observations in the Lower Troposphere Above Kyzyl-Kumami (Aktinometricheskiye nablyudeniya v nizhney troposfere nad Kyzyl-Kumami)

PERIODICAL: Tr. Tsentr. aerolog. observ., 1954, Nr 13, pp 27-37

ABSTRACT: Aircraft-borne actinometric observations were performed above Kyzyl-Kumami and the Mountain District of Turkestan during April and May of 1951. The measurement comprised the intensity of the incoming (downward) and outgoing (upward) fluxes of shortwave radiation, the intensity of the direct and scattered radiation, and the basic meteorological elements. The observations were performed at elevations of 0.5, 1.5, 3.0, and from 4 to 5 km. Twelve flights were carried out in all. Data analysis showed that the density of the downward flux increases at all times with increasing elevation; the vertical gradient of this quantity against elevation, in general, decreases with elevation. The density of the upward flux varies without any definite trend up to the 1.5-km level and

Card 1/2

124-58-9-10068

Actinometric Observations in the Lower Troposphere (cont.)

increases above that level. Both the intensity of the absorption of the insolation and the radiational heating of the air decrease with increasing elevation; the decrease is most pronounced at the 3-km level and near the ground surface. The values of the flux density of the direct and scattered (clear-sky) radiation above Kyzyl-Kumami are close to the values observed at corresponding elevations in the mountainous area. The albedo of the desert surface and of the air layer lying between the aircraft and the ground surface increases with decreasing angular elevation of the sun above the horizon. This, apparently, is governed by the properties of the desert surface but not by those of the inter-jacent air layer. The character of the change in albedo with elevation depends on the ratio between the processes of absorption and scattering of the radiation in the air layer interceding between the aircraft and the ground surface.

L. I. Mamontova

1. Radioactivity--Measurement 2. Atmosphere--Radioactivity

Card 2/2

LASTROV, V.G.; FARAPONOVA, G.P.

Radiation errors of a platinum resistance thermometer. Trudy TSAO
no.16:18-20 '56. (MLRA 9:11)
(Atmospheric temperature) (Thermometers)

~~FARAPONOVA, G.P.~~

Measuring the weakening factor of sunlight in the atmosphere.

Trudy TSAO no.23:52-62 '57.

(MIRA 11:5)

(Atmosphere) (Photometry) (Solar radiation)

~~FARAPONOVA, G.D.~~

Measuring the weakening of sunlight in the free atmosphere.
Trudy TSAO no.32:3-16 '59. (MIRA 12:12)
(Atmospheric transparency)

S/169/61/0
D228/D304

3,5150

AUTHOR:

Faraponova, G.P.

TITLE:

Measuring the weakening of sunlight in the free atmosphere

PERIODICAL:

Referativnyy zhurnal, Geofizika, no. 11, 1961, 26-27.
abstract 11B193 (Tr. Tsentr. aerol. observ., no. 32
1959, 3 - 16)

TEXT: The index of the weakening of sunlight was determined from measurements of direct solar illumination at different altitudes in the atmosphere during aircraft flights to a height of 6.5 km. The observations were conducted with an electrophotometer in which a $\Phi 3Y-19$ (FEU-19) photomultiplier was used as the radiation receiver. Interference filters with passage maxima of 420, 452, 496, and 550 m μ were employed in the equipment. The observations were made in the North Caucasus and in the Khar'kov region in 1957 and in the Moscow region in 1958. On comparing the data obtained in the free atmosphere and in mountainous terrain it was found that the weaken-

Card 1/3

32278

S/169/61/000/011/036/065
D228/D304

Measuring the weakening of ...

ing of light is somewhat greater in the free atmosphere than in the mountains. The optical densities obtained for the free atmosphere also have larger values. It was discovered that the turbidity of the air in the near-surface layer up to 1 km very rapidly diminishes with altitude. A layer with a comparatively constant turbidity from 1 to 3 km is very often observed in summertime; the upper boundary of this layer usually reached the inversion level. Above 3 km the index of weakening declines almost exponentially with altitude. In the lower layers of the troposphere (up to 3 - 4 km) dilution by aerosols plays a leading role in the values of the index of weakening; at greater heights (from 3 - 4 to 6.5 km) the molecular component equals, or becomes somewhat larger than, the aerosol component. In the summer period in the lower layers (up to 4 km) the indices of weakening have somewhat higher values in steppe districts (in North Kazakhstan and the North Caucasus) than in the central part of the Union's European territory (in the Moscow and Khar'kov regions). Observations in the free atmosphere, carried out in West Germany by U. Krug-Pielsticker (Ber. Dtsch. Wetterdienst, no. 8, 1949) and in England by I.M. Waldram (Quart. J. Roy. Meteor.

Card 2/3

S/169/62/000/003/048/098
D228/D301

3,5150

AUTHOR: Faraponova, G. P.

TITLE: The weakening of sunlight in the free atmosphere (Theses)

PERIODICAL: Referativnyy zhurnal, Geofizika, no. 3, 1962, 27; abstract 3B215 (V sb. Aktinometriya i atmosfer. optika, L., Gidrometeoizdat, 1961, 145-146)

TEXT: Measurements were made in an aircraft by means of special photometers at different points in the USSR in 1956-1958. In the troposphere's lower layers from 0 to 1 km the atmospheric turbidity rapidly diminishes. Above 3 - 4 km the index of weakening decreases almost exponentially with altitude, as do the aerosol and the molecular components of the index of weakening, too. The atmosphere's optical densities at a height of 6.5 km exceed the Rayleigh values, indicating considerable turbidity in the troposphere's lower layers. [Abstracter's note: Complete translation.]

Card 1/1

1000

ENCL: 00

NO. 1000 ES

1000

OTHER: 100

1000 1000 1000

ml
Card 2/2

ACCESSION NR: AP4034796

S/0293/64/002/002/0257/0265

AUTHOR: Malkevich, M. S.; Malkov, I. P.; Pakhomova, L. A.; Rozenberg, G. V.;
Faraponova, G. P.

TITLE: Determination of the statistical characteristics of radiation fields over
clouds

SOURCE: Kosmicheskiye issledovaniya, v. 2, no. 2, 1964, 257-265

TOPIC TAGS: meteorology, cloud, atmospheric radiation, radiation field

ABSTRACT: A study has been made of the possibility of applying statistical analysis to fields of outgoing radiation for determining the structure of cloud formations. Computation of the structural parameters of the cloud cover is accomplished using aircraft measurements of radiation with narrow- and wide-angle instruments. The following conclusions are drawn from this preliminary investigation: 1. Statistical characteristics of the intensity of reflected radiation can be used for an objective analysis of clouds of various types and a reliable identification can be made on the basis of the full set of statistical parameters. 2. The most informative parameter is the spectral density of fluctuations of brightness, which is quite sensitive to a difference in the character of nonhomogeneities of different cloud types and at the same time is statistically stable. 3. An investi-

Cord

ACCESSION NR: AP4034796

gation of the statistical characteristics of radiation fluxes, considered as random functions, makes it possible to take into account fluctuations of the radiant flux of heat under conditions of arbitrary cloudiness. In this case spectral density makes it possible to obtain the distribution of radiant energy by frequencies and determine those scales of nonhomogeneities which make the principal contribution to the flux of radiation heat. 4. The spectrum of fluctuations is similar to comparable spectra of fluctuations of wind velocity and temperature obtained in investigations of turbulence in the surface layer of the air. The spectrum was displaced into the region of somewhat lower frequencies, evidence of an increase in the scales of the eddies responsible for the nonhomogeneity of cloud formations. Orig. art. has: 10 formulas, 6 figures and 1 table.

ASSOCIATION: none

SUBMITTED: 23Dec63

SUB CODE: ES

DATE ACQ: 20May64

NO REF SOV: 009

ENCL: 00

OTHER: 003

2/2
Card